

Optimal Renormalization-Group Improvement of $R(s)$ via the Method of Characteristics

V. Elias ^{a,b,1}, D.G.C. McKeon ^{b2} and T. G. Steele ^{c3}

^a *Perimeter Institute for Theoretical Physics, 35 King Street North,
Waterloo, Ontario N2J 2W9 CANADA*

^b *Department of Applied Mathematics, The University of Western Ontario,
London, Ontario N6A 5B7 CANADA*

^c *Department of Physics and Engineering Physics, University of Saskatchewan,
Saskatoon, Saskatchewan S7N 5C6 CANADA*

Abstract

We discuss the application of the method of characteristics to the renormalization-group equation for the perturbative QCD series within the electron-positron annihilation cross-section. We demonstrate how one such renormalization-group improvement of this series is equivalent to a closed-form summation of the first four towers of renormalization-group accessible logarithms to all orders of perturbation theory.

¹Electronic address: velias@uwo.ca

²Electronic address: dgmckeo2@uwo.ca

³Electronic address: steelet@sask.usask.ca

The renormalization group (RG) has played a central role in our understanding of quantum field theory [1-10] especially since the discovery of asymptotic freedom [11-14]. The central idea of the renormalization group is the insensitivity of physical quantities to the mass scale μ^2 introduced in the course of regularizing and eliminating infinities within perturbative calculations. Explicit dependence of a perturbative series on μ^2 is compensated by μ^2 dependence in masses and coupling constants characterising that series. Indeed, the replacement of such quantities by running quantities that are explicitly functions of μ^2 constitutes what is generally denoted by “RG-improvement” of a perturbative expression [15]. The numerical value of a calculation to a given order of perturbation theory still depends upon the numerical value of μ^2 , entailing the introduction of either prescriptions (e.g., $m_b/2 \leq \mu \leq 2m_b$ for semileptonic b -decays) or procedures [16, 17] to obtain optimal values of μ^2 .

However, such substitutions do not in themselves take full advantage of all information accessible from the renormalization-group equation (RGE), which also determines portions of the perturbative series beyond the order of perturbation theory to which calculations have been explicitly performed. Application of the RGE to one-loop expressions has long been known to determine the leading logarithm contribution to each subsequent order of perturbation theory. The RGE can similarly be used in conjunction with two-loop calculations to determine next-to-leading logarithm contributions to all subsequent orders in perturbation theory – indeed the application of the RGE to an n^{th} loop perturbative

expression is sufficient to determine the contribution of n successively-subleading logarithms to all orders in the perturbative expansion parameter.¹ Such RGE methods for obtaining and summing “RG-accessible” logarithms to all orders of perturbation theory have been applied to effective potentials [19] and actions [20, 21], QCD correlation functions [22, 23], QCD contributions to decay rates [22], and even the high-energy behaviour of the $WW \rightarrow ZZ$ cross-section [22], a process dominated by Higgs boson exchanges. A related RG-summation of dimensionality poles in the expansion of the bare coupling constant in terms of its renormalized analog has been developed in ref. [24] and (for thermal field theory) in ref. [25].

The point we wish to emphasize is that the summation of higher order logarithmic contributions is quite distinct (and a substantial improvement over) what is usually understood to be RG-improvement, the incorporation of running masses and coupling constants into perturbative expressions taken to a given order. Indeed, such inclusion of all RG-accessible logarithms within perturbative series is forcefully advocated in ref. [26]. Series which incorporate summation of RG-accessible logarithmic contributions to all orders of perturbation theory have been seen to exhibit much less dependence on μ^2 than series which utilize running masses and coupling constants to a fixed calculational order [22, 23]. This latter approach, however, devolves from the method of characteristics [27], a standard approach to first-order partial differential equa-

¹The possibility that infrared effects might alter this is addressed in ref. [18]

tions such as the RGE [28]. We demonstrate below how this same method of characteristics can be extended to obtain summations of leading and three successively-subleading towers of logarithms to all orders of the perturbative QCD series for the electron-positron annihilation cross-section.

The total cross-section for e^+e^- -annihilation, $R(s) \equiv \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ can be extracted from the imaginary part of the QCD vector-current correlation function [29], a perturbative expression that necessarily depends upon a renormalization mass scale μ :

$$R(s) = 3 \sum_f Q_f^2 S[x(\mu^2), \log(\mu^2/s)] . \quad (1)$$

The expansion parameter $x(\mu^2) \equiv \alpha_s(\mu^2)/\pi$ is proportional to the running QCD coupling constant, and renormalization mass scale μ is a by-product of the regularization procedure for indentifying and excising infinities from the underlying correlation function, as discussed above. Since $R(s)$ cannot depend on this unphysical scale parameter, it follows that

$$\mu^2 \frac{dR(s)}{d\mu^2} = 0 = \left(\mu^2 \frac{\partial}{\partial \mu^2} + \beta(x) \frac{\partial}{\partial x} \right) R(s) \quad (2)$$

where

$$\beta(x(\mu^2)) \equiv \mu^2 \frac{dx(\mu^2)}{d\mu^2} \quad (3)$$

with an appropriately chosen boundary condition for (3) [*e.g.* $x(\Lambda^2) = \infty$ or $x(M_z^2) = 0.118/\pi$]. Perturbative series expansions of $S[x(\mu^2), \log(\frac{\mu^2}{s})]$ and $\beta(x)$ are seen to take the form

$$S[x, L] = 1 + \sum_{n=1}^{\infty} \sum_{m=0}^{n-1} T_{n,m} x^n L^m \quad (4a)$$

$$\beta(x) = -x^2 \sum_{k=0}^{\infty} \beta_k x^k \quad (4b)$$

where $x = x(\mu^2)$ and $L \equiv \log(\mu^2/s)$. Generally $\beta(x)$ is determined by relating the bare coupling to the renormalized coupling [7, 9, 24], although it can also be extracted directly from eq. (2) [20, 21, 30]. Indeed, explicit Feynman diagrammatic calculations to four-loop order have determined $\beta_0, \beta_1, \beta_2, \beta_3$ [13, 14, 31] as well as $T_{1,0}, T_{2,1}, T_{2,0}, T_{3,2}, T_{3,1}$ and $T_{3,0}$ [29], and these results (as tabulated in Table I of ref. [23]) are manifestly consistent with eq. (2).

However, it is possible to utilize eq. (2) to extract higher-order coefficients within $S[x, L]$ than those tabulated in Table 1. It is easily seen [23] that $T_{1,0}$ and β_0 determine *all* leading logarithm coefficients $T_{n,n-1}$ for $n > 1$; similarly additional knowledge of $T_{2,0}$ and β_1 is sufficient to determine all next-to-leading logarithm coefficients $T_{n,n-2}$ for $n > 2$; knowledge of $T_{3,0}$ and β_2 permits determination of $T_{n,n-3}$ for $n > 3$, and so forth. In ref. [23], the double summation in eq. (4a) is reorganised into the form

$$S[x, L] = 1 + \sum_{n=1}^{\infty} x^n S_n(xL) \quad (5)$$

where the functions

$$S_n(u) = \sum_{k=0}^{\infty} T_{n+k,k} u^k \quad (6)$$

are completely determined by knowledge of the “RG-accessible” coefficients

$T_{n+k,k}$. One can show that eq. (2) gives rise to a nested set of first order differential equations for the functions $S_n(u)$:

$$\frac{dS_k}{du} - \frac{k\beta_0}{1-\beta_0 u} S_k = \frac{(1-\delta_{k,1})}{1-\beta_0 u} \sum_{\ell=1}^{k-1} \beta_\ell \left(u \frac{d}{du} + k - \ell \right) S_{k-\ell}(u), \quad S_n(0) = T_{n,0}. \quad (7)$$

These equations are derived and sequentially solved in ref. [23]. When one applies this “RG-summation” to the series (5) within $R(s)$, the dependence of $R(s)$ on μ^2 is substantially reduced [22, 23]. This is not surprising, as the exact result for $R(s)$ is necessarily independent of μ^2 [the RGE is just a statement of this independence], and the inclusion of higher-order logarithm contributions to $R(s)$ via (5) is expected to approximate the exact result more closely than truncation of eq. (4) to a given order.

As discussed above, the method of characteristics [27,28] provides a complementary procedure for obtaining information from the renormalization group equation (2). To illustrate this method, consider the first-order partial differential equation

$$\left[f(x, y) \frac{\partial}{\partial x} + g(x, y) \frac{\partial}{\partial y} \right] A(x, y) = 0 \quad (8)$$

where f and g are given functions, and where $A(x, y)$ may be indentified as some field-theoretical amplitude characterised by quantities (*e.g.*, coupling constants) x and y . If $A_0(x, y)$ is a solution to eq. (8), then so is $A_0(\bar{x}(t), \bar{y}(t))$, provided that

$$\frac{d\bar{x}}{dt} = f(\bar{x}(t), \bar{y}(t)), \quad \frac{d\bar{y}}{dt} = g(\bar{x}(t), \bar{y}(t)) \quad (9)$$

with initial conditions $\bar{x}(0) = x$, $\bar{y}(0) = y$. One then sees from eqs. (9) that

$$0 = \left[f(\bar{x}, \bar{y}) \frac{\partial}{\partial \bar{x}} + g(\bar{x}, \bar{y}) \frac{\partial}{\partial \bar{y}} \right] A_0(\bar{x}, \bar{y}) = \frac{d}{dt} A_0(\bar{x}(t), \bar{y}(t)) \quad (10)$$

The initial conditions ensure that $A_0(\bar{x}(t), \bar{y}(t))$ is a solution of eq. (8) when $t = 0$. Since eq. (10) implies that $A_0(\bar{x}(t), \bar{y}(t))$ is independent of t , $A_0(\bar{x}(t), \bar{y}(t))$ is necessarily a solution to eq. (8) for all values of t . Eqs. (9) and (10) provide the justification for replacing the variables x and y with their corresponding characteristic functions $\bar{x}, \bar{y}$ in the amplitude $A(x, y)$.

For the RGE (2), as applied to the field theoretical series $S[x, \log(\mu^2/s)]$, the role of f and g as dependent variables is assumed by μ^2 and β , in which case correspondence to eqs. (9) requires running values for these functions

$$\frac{d\bar{\mu}^2(t)}{dt} = \bar{\mu}^2(t), \quad (11)$$

$$\frac{d\bar{x}(t)}{dt} = \beta(\bar{x}(t)). \quad (12)$$

The usual prescription for RG improvement is to identify t with $\log(\mu^2)$ [*i.e.*, $\bar{\mu}^2 = e^t = \mu^2$ via eq. (11)], in which case eq. (12) becomes eq. (3). Indeed, this construction provides the justification for having the coupling constant x run with μ^2 within the perturbative series (4a) [27].

However, it is entirely valid to let t be arbitrarily chosen in eqs. (11) and (12), up to initial conditions $\bar{\mu}^2(0) = \mu^2$, $\bar{x}(0) = x(\mu^2)$ that establish contact with a known solution to eq. (2). Thus the “running coupling” $x(\mu^2)$ may be employed to serve as an initial condition for the characteristic function $\bar{x}(t)$.²

²The dimensional regularization equation relating the bare (g_B) and renormalized (g) cou-

In order to keep track of the order of perturbation theory to which we are working, we follow the approach of ref. [32] by introducing an expansion parameter $\hbar$ such that

$$t \longrightarrow t/\hbar \quad (13a)$$

$$\bar{x} \longrightarrow \bar{x}\hbar \quad (13b)$$

so that the characteristic equations (9) become

$$\hbar \frac{d\bar{\mu}^2(t)}{dt} = \bar{\mu}^2(t) \quad (14)$$

$$\hbar^2 \frac{d\bar{x}(t)}{dt} = -\bar{x}^2 \hbar^2 \sum_{n=0}^{\infty} \bar{x}^n \hbar^n \beta_n \quad (15)$$

and the series expansion (4a) becomes

$$S[\bar{x}, \bar{L}] = 1 + \sum_{n=1}^{\infty} \sum_{m=0}^{n-1} T_{n,m} \hbar^n \bar{x}^n \bar{L}^m \quad (16)$$

$[\bar{L} \equiv \log(\bar{\mu}^2(t)/s)]$. From eq. (14) and the $\bar{\mu}(0) = \mu$ initial condition, we see that

$$\bar{\mu}^2(t) = \mu^2 e^{t/\hbar}. \quad (17)$$

We now express $\bar{x}(t)$ as a perturbative expansion

$$\bar{x}(t) = \sum_{n=0}^{\infty} \bar{x}_n(t) \hbar^n \quad (18)$$

pling constants is $g_B = \mu^\epsilon \sum_{\ell=0}^{\infty} \sum_{k=\ell}^{\infty} a_{k,\ell} g^{2k+1}/\epsilon^\ell$ [7, 9]. Since g_B is a bare parameter independent of μ , the renormalized coupling-constant g is necessarily a μ -dependent quantity, *i.e.*, a function of μ .

with $\bar{x}_0(0) = x(\mu^2)$ and $\bar{x}_n(0) = 0$ for $n > 0$. Upon substituting eq. (18) into eq. (15), we obtain a nested set of linear first-order differential equations for the variables $\bar{x}_n(t)$ when $n > 0$:

$$\frac{d\bar{x}_0}{dt} = -\beta_0 \bar{x}_0^2, \quad \bar{x}_0(0) = x(\mu^2), \quad (19)$$

$$\frac{d\bar{x}_1}{dt} + (2\beta_0 \bar{x}_0(t)) \bar{x}_1 = -\beta_1 \bar{x}_0^3(t), \quad \bar{x}_1(0) = 0, \quad (20)$$

$$\frac{d\bar{x}_2}{dt} + (2\beta_0 \bar{x}_0(t)) \bar{x}_2 = -\beta_2 \bar{x}_0^4(t) - 3\beta_1 \bar{x}_0^2(t) \bar{x}_1(t) - \beta_0 \bar{x}_1^2(t), \quad \bar{x}_2(0) = 0, \quad (21)$$

$$\begin{aligned} \frac{d\bar{x}_3}{dt} + (2\beta_0 \bar{x}_0(t)) \bar{x}_3 = & -\beta_3 \bar{x}_0^5(t) - 4\beta_2 \bar{x}_0^3(t) \bar{x}_1(t) \\ & - 3\beta_1 (\bar{x}_0^2(t) \bar{x}_2(t) + \bar{x}_0(t) \bar{x}_1^2(t)) \\ & - 2\beta_0 \bar{x}_1(t) \bar{x}_2(t), \quad \bar{x}_3(0) = 0. \end{aligned} \quad (22)$$

The solution to eq. (19) is

$$\bar{x}_0(t) = \frac{x(\mu^2)}{1 + \beta_0 x(\mu^2)t}. \quad (23)$$

If we substitute eq. (23) into eq. (20), we find the solution to eq. (20) to be

$$\bar{x}_1(t) = -\frac{\beta_1}{\beta_0} \frac{x^2(\mu^2) \log(1 + \beta_0 x(\mu^2)t)}{(1 + \beta_0 x(\mu^2)t)^2}. \quad (24)$$

Similarly, substitution of eq. (23) and (24) into eq. (21) leads to a solution for

$\bar{x}_2(t)$

$$\begin{aligned} \bar{x}_2(t) = & \frac{x^3(\mu^2)}{(1 + \beta_0 x(\mu^2)t)^3} \left[\left(\frac{\beta_1^2}{\beta_0^2} - \frac{\beta_2}{\beta_0} \right) \beta_0 x(\mu^2)t - \frac{\beta_1^2}{\beta_0^2} \log(1 + \beta_0 x(\mu^2)t) \right. \\ & \left. + \frac{\beta_1^2}{\beta_0^2} \log^2(1 + \beta_0 x(\mu^2)t) \right], \end{aligned} \quad (25)$$

and substitution of eqs. (23), (24) and (25) into eq. (22) leads to a solution of

$\bar{x}_3(t)$:

$$\begin{aligned}
\bar{x}_3(t) = & \frac{x^4(\mu^2)}{(1 + \beta_0 x(\mu^2)t)^4} \left[\left(-\frac{\beta_1^3}{2\beta_0^3} + \frac{\beta_1\beta_2}{\beta_0^2} - \frac{\beta_3}{2\beta_0} \right) (1 + \beta_0 x(\mu^2)t)^2 \right. \\
& + \left(\frac{\beta_1^3}{\beta_0^3} - \frac{\beta_1\beta_2}{\beta_0^2} \right) (1 + \beta_0 x(\mu^2)t) (1 - 2\log(1 + \beta_0 x(\mu^2)t)) \\
& + \left(-\frac{\beta_1^3}{2\beta_0^3} + \frac{\beta_3}{2\beta_0} \right) + \left(\frac{2\beta_1^3}{\beta_0^3} - \frac{3\beta_1\beta_2}{\beta_0^2} \right) \log(1 + \beta_0 x(\mu^2)t) \\
& \left. + \frac{5\beta_1^3}{2\beta_0^3} \log^2(1 + \beta_0 x(\mu^2)t) - \frac{\beta_1^3}{\beta_0^3} \log^3(1 + \beta_0 x(\mu^2)t) \right]. \quad (26)
\end{aligned}$$

(Eqs. (23), (24) and (25) also appear in refs. [21,32].) Eqs. (23) - (26) provide an expansion (18) of the solution to (15) that is distinct from the usual perturbative expansion.

If we substitute the series (18) into the expansion (16) for $S[\bar{x}, \bar{L}]$ we find the following solution to the renormalization group equation (2):

$$\begin{aligned}
S[\bar{x}, \bar{L}] = & 1 + \hbar[T_{1,0}\bar{x}_0(t)] \\
& + \hbar^2 [T_{1,0}\bar{x}_1(t) + (T_{2,0} + T_{2,1}\bar{L})\bar{x}_0^2(t)] \\
& + \hbar^3 [T_{1,0}\bar{x}_2(t) + (T_{2,0} + T_{2,1}\bar{L}) (2\bar{x}_0(t)\bar{x}_1(t)) \\
& \quad + (T_{3,0} + T_{3,1}\bar{L} + T_{3,2}\bar{L}^2) \bar{x}_0^3(t)] \\
& + \hbar^4 [T_{1,0}\bar{x}_3(t) + (T_{2,0} + T_{2,1}\bar{L})(\bar{x}_1^2(t) + 2\bar{x}_0(t)\bar{x}_2(t)) \\
& \quad + (T_{3,0} + T_{3,1}\bar{L} + T_{3,2}\bar{L}^2) (3\bar{x}_0^2(t)\bar{x}_1(t)) \\
& \quad + (T_{4,0} + T_{4,1}\bar{L} + T_{4,2}\bar{L}^2 + T_{4,3}\bar{L}^3) \bar{x}_0^4(t)] \\
& + \dots \quad (27)
\end{aligned}$$

Now, if $t = 0$, we find from eq. (17) that $\bar{\mu}^2 = \mu^2$, $\bar{L} = \log(\mu^2/s) = L$. Since

$\bar{x}_0(0) = x(\mu^2)$ and $\bar{x}_k(0) = 0$ for $k \geq 1$, we see that eq. (27) recovers the original expansion (4a) when $\hbar = 1$.

It t is a non-zero constant, the solution (27) provides a means for obtaining all coefficients $T_{n,m}$ with $m \neq 0$ in terms of coefficients $T_{k,0}$. To see this, let $t = \hbar \log k$, in which case we see from eq. (17) that $\bar{\mu}^2 = k\mu^2$, $\bar{L} = \log(k\mu^2/s) = L + \log k$. If we substitute $t = \hbar \log k$ into eqs. (23) - (26) and note that

$$\begin{aligned} \log(1 + \hbar\beta_0 x(\mu^2)t) &= \hbar\beta_0 x(\mu^2) \log k \\ &- \frac{\hbar^2}{2} \beta_0^2 x^2(\mu^2) \log^2 k \\ &+ \frac{\hbar^3}{3} \beta_0^3 x^3(\mu^2) \log^3 k + \dots, \end{aligned} \quad (28)$$

we find upon further substitution into eq. (27) that

$$\begin{aligned} S[\bar{x}, \bar{L}]|_{t=\hbar \log k} &= 1 + \hbar x(\mu^2) T_{1,0} \\ &+ \hbar^2 x^2(\mu^2) [(T_{2,0} - \beta_0 T_{1,0} \log k) + T_{2,1} \log(k\mu^2/s)] \\ &+ \hbar^3 x^3(\mu^2) \{ [T_{3,0} - (2T_{2,0}\beta_0 + T_{1,0}\beta_1) \log k + T_{1,0}\beta_0^2 \log^2 k] \\ &\quad + [T_{3,1} - 2\beta_0 T_{2,1} \log k] \log(k\mu^2/s) + T_{3,2} \log^2(k\mu^2/s) \} \\ &+ \hbar^4 x^4(\mu^2) \{ [T_{4,0} - (3\beta_0 T_{3,0} + 2\beta_1 T_{2,0} + \beta_2 T_{1,0}) \log k \\ &\quad + \left(3\beta_0^2 T_{2,0} + \frac{5}{2}\beta_0\beta_1 T_{1,0} \right) \log^2 k - \beta_0^3 T_{1,0} \log^3 k] \\ &\quad + [T_{4,1} - (3\beta_0 T_{3,1} + 2\beta_1 T_{2,1}) \log k + 3\beta_0^2 T_{2,1} \log^2 k] \log(k\mu^2/s) \\ &\quad + [T_{4,2} - 3\beta_0 T_{3,2} \log k] \log^2(k\mu^2/s) + T_{4,3} \log^3(k\mu^2/s) \} \\ &+ \mathcal{O}(\hbar^5 x^5(\mu^2)). \end{aligned} \quad (29)$$

Now if we rewrite the original series expansion (4a) with $L \equiv \log(\mu^2/s) =$

$\log(k\mu^2/s) - \log k$, we obtain

$$\begin{aligned}
S[x, L] &= 1 + x(\mu^2)T_{1,0} \\
&+ x^2(\mu^2) [(T_{2,0} - T_{2,1} \log k) + T_{2,1} \log(k\mu^2/s)] \\
&+ x^3(\mu^2) [(T_{3,0} - T_{3,1} \log k + T_{3,2} \log^2 k) \\
&\quad + (T_{3,1} - 2T_{3,2} \log k) \log(k\mu^2/s) \\
&\quad + T_{3,2} \log^2(k\mu^2/s)] \\
&+ x^4(\mu^2) [(T_{4,0} - T_{4,1} \log k + T_{4,2} \log^2 k - T_{4,3} \log^3 k) \\
&\quad + (T_{4,1} - 2T_{4,2} \log k + 3T_{4,3} \log^2 k) \log(k\mu^2/s) \\
&\quad + (T_{4,2} - 3T_{4,3} \log k) \log^2(k\mu^2/s) + T_{4,3} \log^3(k\mu^2/s)] \\
&+ \mathcal{O}(x^5(\mu^2)). \tag{30}
\end{aligned}$$

When $\hbar = 1$, eqs. (29) and (30) must be equal, since the solution (27) to the renormalization-group equation (2) has been constructed...

- 1) ...to coincide with eq. (30) at $t = 0$,
- 2) ...to be independent of the choice for t via the method of characteristics [eq. (10)], and
- 3) ...to be given by eq. (29) when $t = \hbar \log k$.

Direct comparison of eqs. (29) and (30) when $\hbar = 1$ shows that

$$\begin{aligned}
T_{2,1} &= \beta_0 T_{1,0}, \\
T_{3,1} &= 2T_{2,0}\beta_0 + \beta_1 T_{1,0},
\end{aligned}$$

$$\begin{aligned}
T_{3,2} &= \beta_0^2 T_{1,0}, \\
T_{4,1} &= 3\beta_0 T_{3,0} + 2\beta_1 T_{2,0} + \beta_2 T_{1,0}, \\
T_{4,2} &= 3\beta_0^2 T_{2,0} + \frac{5}{2}\beta_0\beta_1 T_{1,0}, \\
T_{4,3} &= \beta_0^3 T_{1,0},
\end{aligned} \tag{31}$$

relations that can also be obtained [22] by direct substitution of the series (4a) into the renormalization-group equation (2). Thus, the method of characteristics is seen to determine all logarithmic coefficients to the order of perturbation theory considered.

However, a more powerful application of the solution (27) occurs by setting $t = \hbar \log(s/\mu^2)$, ensuring via eq. (17) that $\bar{\mu}^2 = s$, that $\bar{L} = 0$, and that factors of $1 + \hbar\beta_0 x(\mu^2)t$ in eqs. (23) - (26) become $1 - \beta_0 x(\mu^2) \log(\mu^2/s) \equiv w$ in the $\hbar \rightarrow 1$ limit. In this limit, eq. (27) generates the following series:

$$\begin{aligned}
S[\bar{x}, \bar{L}] &= 1 + x(\mu^2) \frac{T_{1,0}}{w} + x^2(\mu^2) \left[T_{2,0} - \frac{\beta_1}{\beta_0} T_{1,0} \log w \right] w^{-2} \\
&+ x^3(\mu^2) \left[T_{3,0} - 2T_{2,0} \frac{\beta_1}{\beta_0} \log w + \left(\frac{\beta_1^2}{\beta_0^2} - \frac{\beta_2}{\beta_0} \right) (w-1) - \frac{\beta_1^2}{\beta_0^2} \log w + \frac{\beta_1^2}{\beta_0^2} \log^2 w \right] w^{-3} \\
&+ x^4(\mu^2) \left[\left(-\frac{\beta_1^3}{2\beta_0^3} + \frac{\beta_1\beta_2}{\beta_0^2} - \frac{\beta_3}{2\beta_0} \right) w^2 + \left(\frac{\beta_1^3}{\beta_0^3} - \frac{\beta_1\beta_2}{\beta_0^2} + 2T_{2,0} \left(\frac{\beta_1^2}{\beta_0^2} - \frac{\beta_2}{\beta_0} \right) \right) w \right. \\
&+ \left(-\frac{2\beta_1^3}{\beta_0^3} + \frac{2\beta_1\beta_2}{\beta_0^2} \right) w \log w + \left(-\frac{\beta_1^3}{2\beta_0^3} + \frac{\beta_3}{2\beta_0} - 2T_{2,0} \left(\frac{\beta_1^2}{\beta_0^2} - \frac{\beta_2}{\beta_0} \right) + T_{4,0} \right) \\
&+ \left(\frac{2\beta_1^3}{\beta_0^3} - \frac{3\beta_1\beta_2}{\beta_0^2} - 2T_{2,0} \frac{\beta_1^2}{\beta_0^2} - 3T_{3,0} \frac{\beta_1}{\beta_0} \right) \log w \\
&+ \left(\frac{5\beta_1^3}{2\beta_0^3} + 3T_{2,0} \frac{\beta_1^2}{\beta_0^2} \right) \log^2 w \\
&\left. - \frac{\beta_1^3}{\beta_0^3} \log^3 w \right] w^{-4} + \dots
\end{aligned} \tag{32}$$

This series explicitly reproduces the series (5) obtained via the successive solu-

tions to the differential equations (7). The coefficient functions $S_1(xL)$, $S_2(xL)$, $S_3(xL)$ and $S_4(xL)$, as calculated in ref. [23], are reproduced in eq. (32), demonstrating how the method-of-characteristics approach to the renormalization group equation can recover the results obtained via summation of leading, next-to-leading, and successively subleading logarithm factors to *all* orders of perturbation theory. Such results as eq. (32) represent the optimal possible RG-improvement of the original perturbative series (4a), insofar as they incorporate all RG-accessible coefficients of logarithms occurring within that series.

As a final note, the ambiguity in the choice of k such that $t = \hbar \log k$ [thereby leading to the series of eq. (29)] is equivalent to the ambiguity noted in ref. [25]. In [25], this ambiguity was viewed as a consequence of shifting the initial condition of eq. (7) to the equation (6) defining $S_n(u)$; by replacing $\log(\mu^2/s)$ with $\log(k\mu^2/s) - \log(k)$ in the series (4a), that series becomes

$$S[x, L] = 1 + \sum_{n=1}^{\infty} \sum_{m=0}^{n-1} T'_{n,m} x^n (L')^m \quad (33)$$

where $L' \equiv \log(k\mu^2/s)$ and where $T'_{1,0} = T_{1,0} = 1$, $T'_{2,1} = T_{2,1}$, $T'_{2,0} = T_{2,0} - T_{2,1} \log(k)$, etc. The initial condition in eq. (7) is now replaced by $S_n(0) = T'_{n,0}$. Of course, when one sums to all orders in perturbation theory, the dependence on k within eq. (29) will drop out.

Acknowledgements

We are grateful for research support from the Natural Sciences and Engineering Research Council of Canada. Discussions with A. Rebhan were most helpful.

V. E. would like to thank R. and D. MacKenzie for discussion.

References

- [1] E. C. G. Stueckelberg and A. Peterman, *Helv. Phys. Acta* 26 (1953) 499.
- [2] N. W. Bogoliubov and D. V. Shirkov, *Introduction to the Theory of Quantized Fields* (John Wiley, New York, 1980) Chapter 9.
- [3] M. Gell-Mann and F. Low, *Phys. Rev.* 95 (1954) 1300.
- [4] K. G. Wilson, *Phys. Rev. D* 3 (1971) 1818.
- [5] C. G. Callan, *Phys. Rev. D* 2 (1970) 1541.
- [6] K. Symanzik, *Commun. Math. Phys.* 18 (1970) 227.
- [7] G. 't Hooft, *Nucl. Phys. B* 61 (1973) 455.
- [8] S. Weinberg, *Phys. Rev. D* 8 (1973) 3497.
- [9] J. C. Collins and A. J. Macfarlane, *Phys. Rev. D* 10 (1974) 1201.
- [10] P. M. Stevenson, *Ann. of Phys.* 132 (1981) 383.
- [11] V. S. Vanyashin and M. V. Terentev, *JETP* 21 (1965) 375.
- [12] I. B. Khriplovich, *Sov. J. Nucl. Phys.* 10 (1970) 235.
- [13] D. J. Gross and F. Wilczek, *Phys. Rev. Lett.* 30 (1973) 1343.
- [14] H. D. Politzer, *Phys. Rev. Lett.* 30 (1973) 1346.

- [15] S. Coleman and E. Weinberg, Phys. Rev. D 7 (1973) 1888.
- [16] P. M. Stevenson, Phys. Rev. D 23 (1981) 2916.
- [17] G. Grunberg, Phys. Lett. B 95 (1980) 70 and Phys. Rev. D 29 (1984) 2315.
- [18] G. Dunne, H. Geis and C. Schubert, JHEP 0211 (2002) 032.
- [19] B. Kastening, Phys. Lett. B 283 (1992) 287; M. Bando, T. Kugo, N. Maskawa and H. Nakano, Phys. Lett. B 301 (1993) 83; C. Ford, Phys. Rev. D 50 (1994) 7531.
- [20] D. G. C. McKeon, Int. J. Mod. Phys. 37 (1998) 817.
- [21] M. R. Ahmady, V. Elias, D. G. C. McKeon, A. Squires and T. G. Steele, Nucl. Phys. B (to appear) hep-ph/0211227.
- [22] M. R. Ahmady *et. al.*, Phys. Rev. D 66 (2002) 014010.
- [23] M. R. Ahmady *et. al.*, Phys. Rev. D (to appear) hep-ph/0208025.
- [24] V. Elias and D. G. C. McKeon, Mod. Phys. Lett. A (to appear) hep-th/0209151.
- [25] A. Rebhan and D. G. C. McKeon, Phys. Rev. D (to appear) hep-ph/0210163.
- [26] C. J. Maxwell, Nucl. Phys. B (Proc. Suppl.) 86 (2000) 74.
- [27] A. Peterman, Phys. Rep. 53 C (1979) 157.

- [28] R. Courant and D. Hilbert, *Methods of Mathematical Physics Vol. II* (Interscience, NY, 1966) Ch. II.
- [29] K. G. Chetyrkin, Phys. Lett. B 391 (1997) 402; see also S. G. Gorishny, A. L. Kataev and S. A. Larin, Phys. Lett. B 259 (1991) 144, and L. R. Surguladze and M. A. Samuel, Phys. Rev. Lett. 66 (1991) 560 and 2416 (E).
- [30] L. Culumovic, M. Leblanc, R. B. Mann, D. G. C. McKeon and T. N. Sherry, Phys. Rev. D 41 (1990) 514; L. Culumovic, D. G. C. McKeon and T. N. Sherry, Ann. Phys. 197 (1990) 94.
- [31] W. E. Caswell, Phys. Rev. Lett. 33 (1974) 244; D. R. T. Jones, Nucl. Phys. B 75 (1974) 531; E. S. Egorian, O. V. Tarasov, Theor. Mat. Fiz. 41 (1979) 26; O. V. Tarasov, A. A. Vladimirov and A. Yu. Zharkov, Phys. Lett. B 93 (1980) 429; S. A. Larin and J. A. M. Vermaseren, Phys. Lett. B 303 (1993) 334; T. van Ritbergen, J. A. M. Vermaseren and S. A. Larin, Phys. Lett. B 400 (1997) 379.
- [32] J. M. Chung and B. K. Chung, Phys. Rev. D 60 (1999) 105001.